

Seyed Mostafa Moshirian
Farahi ¹

Alireza Iranbakhsh ²

Homa Mahmoodzadeh ³

Mostafa Ebadi ⁴

Javad Baharara ³

Effects of titanium dioxide nanoparticles (TiO₂) on germination and seedling growth of Vitex plants (*Vitex agnus-castus* L.)

Authors' addresses:

¹ Department of Biology, Islamic Azad University, Science and Research Branch, Tehran, Iran.

² Department of Biology, Science Faculty, Science and Research Azad University, Iran.

³ Department of Biology, Islamic Azad University of Mashhad, Khorasan Razavi Province, Iran

⁴ Department of Biology, Islamic Azad University of Damghan, Semnan Province, Iran

Correspondence:

Alireza Iranbakhsh

Department of Biology, Science Faculty, Science and Research Azad University, Daneshgah Sq., Sattari Hwy, Tehran, Iran.

Tel.: +989121487477

e-mail: iranbakhshar@yahoo.com

Article info:

Received: 16 February 2020

Accepted: 1 April 2020

ABSTRACT

Different concentrations of nanoparticles impose impact different effects on germination and plant growth. The purpose of this study was to investigate the effects of 11 different concentrations of titanium dioxide nanoparticles on germination and seedling growth characteristics of Vitex plant (*Vitex agnus-castus* L.). This experiment was conducted in the laboratory of the Islamic Azad University of Tehran in 2018 as a completely randomized design with 4 replications. The measured traits included germination percentage, relative germination percentage, germination rate, mean germination time, germination index, weighted germination index, vigor seedling index, fresh and dry weight of seedling and root and shoot length. XRD measurements showed that the titanium dioxide nanoparticles used in the anatase form were present. The results showed that the highest germination percentage (80%), relative germination percentage (115%), germination rate (22.37), germination index (190), weighted germination index (2.30), vigor seedling index (541) and fresh weight (0.52) were observed through application of 200 µg.ml⁻¹ titanium nitric oxide. Higher concentrations of nano titanium dioxide had no positive effect on germination indices. The response index also showed that nano-titanium dioxide at all concentrations had a stimulatory effect on mean germination time, root length, shoot length and dry weight but only at 200 mg.l⁻¹ treatment had a stimulatory effect on germination percentage and germination rate. The results of this study indicated that the treatment of 200 µg.ml⁻¹ titanium dioxide nanoparticles had the most positive effect on germination and seedling growth factors of Vitex plantlets.

Key words: Nanoparticles, Vitex (*Vitex agnus-castus* L.), Titanium, Germination indices, Response index

Introduction

Nanotechnology is one of the most extensive and attractive sciences that has many applications in various fields such as agriculture, industry, biotechnology, electronics, medicine, energy, and life sciences. (Nair & Chung, 2014; Tripathi *et al.*, 2017; Tiwari *et al.*, 2019). Research on nanoparticles is increasing rapidly due to distinct physical and chemical properties such as particle morphology, larger surface area, pore size, high reactivity, and physical and chemical properties related to particle size (Siddiqui *et al.*, 2015). Today, large amounts of nanoparticles for use in various disciplines and industry research have led to an increase in their applications. These nanoparticles and their intermediate compounds interact with plants, animals,

and the other living systems and expose positive and negative effects (Tripathi *et al.*, 2017). Most nanoparticles have been reported to have toxic effects even at very low concentrations and affect the morphophysiological, biochemical, and molecular properties of living systems. However, some nanoparticles also have positive effects on plants by increasing their growth and protecting them against various stresses (Syu *et al.*, 2014; Rico *et al.*, 2015; Tripathi *et al.*, 2016). In recent years, the use of titanium dioxide nanoparticles due to their biological properties has attracted much attention from plant physiologists (Qi *et al.*, 2013). Dioxide nanoparticles have high soluble cyanides and can serve as water-soluble carriers, iron oxides, ammonium salts, and energy-soluble kinases (Kenanakis & Katsarakis, 2014). TiO₂ nanoparticles are available in three crystalline forms of

anatase (a quadrilateral with a density of 0.8 g.cm^{-3}), rutile (a quadrilateral with a density of 2.9 g.cm^{-3}) and brookite (amorphous with a density of 0.5 g.cm^{-3}) (Afzali & Maghsoodlou, 2016). Titanium is the ninth most abundant element in the earth's crust, albeit secondarily in a transition metal, and evidence suggests that it can increase crop yields by about 10 to 20 percent (Feizi *et al.*, 2013). It can also stimulate the absorption of some elements such as nitrogen, phosphorus, calcium, magnesium, iron, manganese and zinc (Marschner, 2011). Nano-TiO₂ is capable of enhancing plant fresh and dry weight by improving light absorption and activity of Rubisco (Mingyu *et al.*, 2007) as well as by increasing nitrate uptake and accelerating the conversion of inorganic matter (Nair *et al.*, 2010). Du *et al.* (2011) showed that due to nanoparticles the growth rate of wheat (*Triticum aestivum*) roots were more than shoots, also compared to control, both roots and shoots showed significantly higher growth. Jaberzadeh *et al.* (2013) reported that Nano titanium dioxide improved wheat growth and its yield components under drought stress. Combination of TiO₂ and SiO₂ nanoparticles in mungbean (*Vigna radiate*) and chickpea (*Cicer arietinum*) increased nitrate reductase activity, germination and growth (Mahajan *et al.*, 2011). Studies have shown that germination of tomato (*Solanum lycopersicum*), lettuce (*Lactuca sativa*) and onion (*Allium cepa* L.) seeds increased in the presence of TiO₂ and after exposure to light (Elghniji *et al.*, 2014). Despite research on crops, there is still limited information on the effects and physiological changes induced by these nanoparticles on germination and growth of plants, especially medicinal and forest plants.

Vitex is the largest genus in the Lamiaceae family, comprising of 250 species worldwide (Rani & Sharma, 2013). Species of this genus are shrubs. One of the most important species used in medicine is the Vitex or Chaste tree (*Vitex agnus-castus* L.) and grown in natural ecosystems in the Mediterranean regions of southern Europe and Central Asia by the riversides and along the waterways (Rani & Sharma, 2013). The fruit of this plant is used to treat menstrual problems in women, including premenstrual syndrome, disruption of estrogen and progesterone hormone balance, menopause, menopausal complications, hyperprolactinemia (Carmichael, 2008; Dugoua *et al.*, 2008). In Iranian traditional medicine, its leaves and fruits are also used to increase human milk production (Azadbakht *et al.*, 2005). One of the major problems in the cultivation and operation of the Vitex is the lack of proper germination and consequently lack of proper establishment in the field conditions. Therefore, it is necessary to apply treatments to improve its germination. The effect of nanoparticles on Vitex seed germination indices has not been studied so far, and due to applications and use of nanoparticles in agronomy and the end-production and cultivation of different crops, it is the

goals of this study to investigate the effects of different concentrations of nanoparticles on germination characteristics of Vitex seeds.

Materials and Methods

Plant Material and Treatment Levels

Vitex seeds were obtained from Dashtyar Isfahan Company in 2018. Seeds were then sterilized with 7% sodium hypochlorite for 10-15 minutes and rinsed three times with distilled water. Then, 25 seeds were located in each 10 cm diameter Petri-sterile. After adding 10 ml of each treatment solution to the Petri, the prepared petri was transferred to a germinator at $25 \pm 1^\circ\text{C}$. Seeds were counted on a daily basis. The germination criterion was when roots of 2 mm length. Distilled water used as control and concentrations of titanium dioxide nanoparticles of 10, 100, 200, 300, 400, 500, 600, 700, 800, 900 and 1000 mg.l^{-1} titanium dioxide nanoparticles were applied as treatments with four replications. Titanium dioxide nanoparticles were obtained from Nutirino Co., Tehran. To provide a uniform solution of titanium dioxide, in addition to the usual shaker, the solution was placed in an ultrasonic bath for 30 minutes and then added to the Petri dishes.

Study factors

Characterization of nanoparticles including spectroscopy, purity and particle size was performed by nanoparticle sizer and XRD device at Mashhad Central Laboratory, Ferdowsi University of Mashhad. At the end of the experiment, after measuring the root and shoot length by the ruler, the weights of the plant parts were measured with digital scales at the accuracy level of mg. These organs were then placed in an oven at 72°C for 48 hours and weighed to determine the dry weight (Yang *et al.*, 2007). The following equations were used to calculate germination parameters.

Equation 1) Germination percentage

$$GP = \frac{GN}{SN} \times 100$$

Where *GN* is the total number of germinated seed on the last day of counting and *SN* is the total number of germinated seed (Panwar & Bhardwaj, 2005).

Equation 2) Relative germination percentage

$$RGP = \frac{GP(\text{treatment})}{GP(\text{control})} \times 100$$

Where *GP* (treatment) is the germination percentage of the treatment and *GP* (control) the germination percentage (Panwar & Bhardwaj, 2005).

Equation 3) Germination rate

$$GR = \sum_n^i \frac{Si}{Gi}$$

Where Si is the number of germinated seeds per desired day and Gi is the number of days elapsed since the start of the experiment (Panwar & Bhardwaj, 2005).

Equation 4) Mean germination time

$$MGT = \sum_i \frac{Gi \times i}{\sum_i Gi}$$

Where Gi is the number of germinated seeds per day and i is the number of days since planting (Kulkarni *et al.*, 2007).

Equation 5) germination index

$$GI = \frac{(\sum(N - i) \times Gi) \times 100}{N \times GN}$$

Where Gi is the number of germinated seeds on day i , i the number of days sown, N the duration of the experiment or the number of study days, and GN is the total number of germinated seeds (Wu & Du, 2007).

Equation 6) Weighted germination index

$$WGI = \frac{[(N \times n1 + (N - 1) \times n2 + (N - 2) \times n3 + \dots)]}{N \times N'}$$

Where N is the total number of study days, N' is the total number of seeds, $n1$, $n2$, $n3$, etc. The number of germinated seeds on the first, second, third day, etc. (Wu & Du, 2007).

Equation 7) Vigour Seedling Index

(Kandil *et al.*, 2015)

$$SVI = GP \times \text{Seedling Length (Root + Shoot)}$$

Equation 8) Optimal Seedling Index

(Vashisth & Nagarajan, 2010)

$$SOI = GP \times \text{Dry Weight}$$

Equation 9) Response Index

(Kandil *et al.*, 2015) If control (C) exceeds treatment (T) then

$$RI = \frac{T}{C} - 1$$

If treatment (T) is greater than control (C) then

$$RI = 1 - C/T$$

Treatments RL (root length), PL (shoot length) and WW (fresh weight of seedling) and DW (dry weight of seedling) was measured.

Statistical analysis was performed using SPSS software version 16 for analysis of variance with the assumption of normality of data. Comparisons were made by Duncan's multiple range test at 5% probability level.

Results

Characterization of titanium dioxide nanoparticles

The crystalline properties of titanium dioxide nanoparticles were investigated by X-ray diffraction (XRD) and the results are presented in (Fig. 1). XRD measurements showed that the titanium dioxide nanoparticles used are in the form of anatase (Fig. 1). According to the size of titanium dioxide nanoparticles which was measured by Nanoparticle size analyzer and shown in (Fig. 2), mean diameter of nanoparticles was 98.87, (D mean), particle hydrodynamic




Figure 1. XRD pattern of TiO₂ nanoparticles.

diameter: 197.01 (Zaverage), and particle dispersion index was 0.2050 (PDI).

Influence of titanium dioxide nanoparticles on germination indices

Analysis of variance showed that treatment with titanium dioxide nanoparticles showed statistically significant changes in most germination and seedling growth traits and the results are summarized in (Table 1).

A comparison of mean germination percentage at different concentrations of titanium dioxide nanoparticles showed that only 100 and 200 $\mu\text{g.ml}^{-1}$ nanoparticles applied, increased germination percentage of Vitex. The highest germination percentage (80%) was obtained from 200 $\mu\text{g.ml}^{-1}$ titanium dioxide treatment, which was 15% more than control (Table 2).

However, higher concentrations decreased germination percentage and the lowest germination percentage was observed in 800, 900 and 1000 $\mu\text{g.ml}^{-1}$ titanium dioxide treatments which decreased by 42, 51 and 51% compared to control, respectively (Table 2).

A comparison of the mean relative germination percentage at different levels of titanium dioxide showed that 100 and 200 $\mu\text{g.ml}^{-1}$ treatment showed the highest relative germination percentage (80%) and relative germination percentage increased by 65% (Table 2). The lowest relative germination percentage (50%) was observed at concentrations of 900 and 1000 $\mu\text{g.ml}^{-1}$ (Table 2).

RESEARCH ARTICLE

Table 1. Results of ANOVA tests for the effects of TiO₂ NPs different concentrations on (*Vitex agnus-castus L.*) studied parameters.

GP	RGP	MGT	GR	GI	WGI	SVI	RL	PL	WW	DW	SOI
.00	.001	.584	.021	.046	.039	.006	.008	.995	.01	.041	0.04

GP: Germination percentage, RGP: Relative germination percentage, MGT: Mean germination time, GR: Germination rate, GI: Germination index, WGI: Weighted germination index, SVI: Vigor seedling index, RL: Root length, PL: Shoot length, WW: Seedling fresh weight, DW: Seedling dry weight, SOI: Optimal Seedling Index.

Table 2. Mean value of Germination Percent (GP%), Relative Germination Percent (RGP%), Mean Germination Time (MGT), Germination Rate (GR), Germination Index (GI) and Weighted Germination Index (WGI) for seeds of (*Vitex agnus-castus L.*) under TiO₂ NPs different concentrations.

TiO ₂ Concentration (ppm)	GP%	RGP%	GR (N/day)	MGT (day)	GI (day)	WGI
Control	69.33ab	69.33c	17.48ab	64.19ab	145ab	1.78ab
10	36.00c	51.91c	8.17bc	68.60ab	76.85bc	0.94bc
100	68.00ab	98.07ab	16.39ab	69.89ab	149ab	1.84ab
200	80.00a	115a	22.37a	66.23ab	190a	2.30a
300	52.00bc	75.00c	10.74bc	65.54ab	96.38bc	1.20bc
400	52.00bc	75.00bc	10.92bc	65.96ab	100bc	1.25bc
500	52.00bc	75.00bc	15.28abc	70.50ab	132abc	1.61abc
600	48.00bc	69.23c	6.30c	65.56b	54.85c	0.72c
700	48.00bc	69.23c	10.83bc	69.77ab	98.47bc	1.22bc
800	40.00c	57.69c	9.84bc	66.74ab	84.90bc	1.04bc
900	34.00c	50.00c	9.20bc	72.29a	83.90bc	1.02bc
1000	34.00c	50.00c	7.77bc	68.90ab	72.19bc	0.89bc

Means followed by different letter(s) show significant difference at (P<0.05) significance level according to the Duncan's multiple ranges test

Treatments that could have a higher germination percentage in less time would also have a higher germination rate. A comparison of the mean germination rate showed that different levels of TiO₂ had different effects on the germination rate. Some treatments decreased the germination rate and some increased it compared to control. The lowest germination rate (3.6 seeds/day) was obtained in 600 µg.ml⁻¹ treatment, which was 63% lower than control (Table 2). The highest germination rate was also observed in 200 µg.ml⁻¹ treatment with an increase of 27% compared to control (Table 2). The positive effect of Nano titanium dioxide on seed germination rate has also been reported in other studies (Feizi *et al.*, 2013).

Undoubtedly, with the introduction of nanoparticles into the seeds, permeability, water and oxygen entry pathway would increase as well therefore germination changes would be expected (Fathi *et al.*, 2017). The positive effects of titanium dioxide on the percentage and rate of germination of sage (Hatami *et al.*, 2014), bitter myrtle (Hatami *et al.*, 2014), onion (Laware & Raskar, 2014) and fennel (Feizi *et al.*, 2013) have also been reported. Titanium dioxide improves seed germination by increasing seed water uptake and oxygen uptake (Zheng *et al.*, 2005). As Nano-titanium particles enter the seed cells, the redox oxidation reactions are intensified by superoxide ion radicals, leading to the release of free radicals into the germinating seeds, as well as the oxygen produced in such a young process. (Zheng *et al.*, 2005; Pais, 2008). Some

studies have found that improving the activity of nitrate reductase and other antioxidants is an effective factor in improving the germination properties of Nano-titanium dioxide-treated seeds (Lu *et al.*, 2002). Clement *et al.* (2013) showed that soaking the seeds in anatase titanium oxide nanoparticles at 100 mg.L⁻¹ had a positive effect on germination and root growth of flax plants. These positive effects could be due to the antimicrobial properties of the TiO₂ crystal structure, which increases the plant's resistance to stress. However, there are also reports pointing at adverse effects of titanium nitric oxide on the germination of some seeds such as tobacco (Frazier *et al.*, 2014) and grass (Azimi *et al.*, 2014). In this experiment, high concentrations of TiO₂ decreased germination indices. This result indicated that as the concentration of nanoparticles increased, their uptake by the seed coat probably increased and caused toxicity in the seed, which decreased with water absorption and decreased germination. Yang and Watts (2005) reported that the effect of Nano titanium dioxide on studied plants (radish, ryegrass, lettuce, maize and cucumber) caused toxicity in the plant and even inhibited root growth. The effects of nanocomposites on plants appear to be different depending on the plant type, growth stage, type of nanoparticle, physicochemical properties of the particle and its amount under different conditions (Aghdam *et al.*, 2016). There was no statistically significant difference between the mean of different levels of titanium dioxide for the mean germination time (Table 1).




Figure 2. Size of titanium dioxide nanoparticles measured by nanoparticle sizer with: a) scattered intensity, b) scattered volume, and c) scattered number

The highest germination time was obtained at a concentration of 900 $\mu\text{g.ml}^{-1}$, which was 8% more than control (Table 2).

In general, the germination index is one of the important parameters in determining seed germination and is directly related to quality and bioenergy of seeds.

A comparison of the mean germination index showed that among different levels of titanium nitric oxide, the germination index increased only at two concentrations of 100 and 200 $\mu\text{g.ml}^{-1}$, (Table 2). The highest germination index of 190 was obtained under 200 $\mu\text{g.ml}^{-1}$ with 31% higher compared to control.

The maximum germination index was observed under 600 $\mu\text{g.ml}^{-1}$ treatment. The maximum and minimum were 2.3 and 0.72 under concentrations of 200 and 600 $\mu\text{g.ml}^{-1}$, respectively. It should be noted that the weighted germination index also increased only at concentrations of 100 and 200 $\mu\text{g.ml}^{-1}$ compared to the control treatment. The vigor seedling index represents the germination percentage and germination potential. This index is one of the most important traits whose increase is a sign of increased quality and germination energy of the seed. Quantitatively this trait is the product of seedling length by germination percentage therefore and treatment which increases seedling length and germination percentage is able to enhance the seeds vigor. The lowest value of Vigour Seedling Index was obtained at concentrations of 900 and 1000 $\mu\text{g.ml}^{-1}$ which showed a 20% decrease compared to the control (Table 3).

The highest value of this index was observed at 200 $\mu\text{g.ml}^{-1}$ with an increase of 186% compared to control (Table 3). Zhang *et al.* (2008) examined the effect of Nano titanium dioxide on spinach seed germination and showed that Nano-particle treatment significantly increased the germination index and seedling vigor index. Feizi *et al.* (2013) suggested that increasing the vigor index of the seed was beneficial and could improve seedling establishment by nanoparticles, which is due to biological activity and biological mobility of nanoparticles influenced by factors such as their size, shape, chemical nature, stability and bioavailability.

Impact of Titanium Dioxide Nanoparticles on Seedling Growth and Biomass

Titanium is a beneficial element for plants and stimulates their growth (Nair *et al.*, 2010). The subsequent effects of these nanoparticles on seedling growth and biomass were also investigated. By comparison, the lowest root length (3.3 mm) was obtained in control plants and the highest root length (26.0 mm) was obtained in 10 $\mu\text{g.ml}^{-1}$ titanium dioxide treatment (Table 3). There was no statistically significant difference in shoot length between different concentrations of nano titanium dioxide (Table 1). The shortest shoot length (24.32 mm) was observed in control and the longest shoot length (36.6 mm) was observed in 1000 $\mu\text{g.ml}^{-1}$ titanium dioxide (Table 3).

A comparison of the mean seedling fresh weight showed that the highest seedling fresh weight as 0.53 mg was observed under 200 $\mu\text{g.ml}^{-1}$ Nano titanium dioxide, which was 32% more than control (Table 3). The lowest seedling fresh weight (0.17 mg) was obtained under 600 $\mu\text{g.ml}^{-1}$ treatment, which was 57% lower than the control (Table 3). According to the analysis of variance, no significant differences were observed between seed treatments for the seedling dry weight (Table 1). The highest seedling dry weight (0.5 $\mu\text{g.ml}^{-1}$) was observed in 1000 $\mu\text{g.ml}^{-1}$ titanium dioxide and the lowest was observed in the control treatment

RESEARCH ARTICLE

Table 3. Mean value of Seedling Vigor Index (SVI), Radicle Length (RL), Plumule Length (PL), Wet Weight (WW) and Dry Weight (DW), Seedling Optimum Index (SOI) for Seedling of (*Vitex agnus-castus L.*) under TiO₂ NPs different concentrations.

TiO ₂ Concentration (ppm)	SVI	RL(mm)	PL(mm)	WW(mg)	DW(mg)	SOI
Control	189 ^{bc}	3.66 ^d	24.33 ^a	0.406 ^{abc}	0.0087 ^d	1.51 ^a
10	196 ^{bc}	26.66 ^a	26.33 ^a	0.333 ^{bcd}	0.0140 ^{cd}	0.86 ^a
100	340 ^{bc}	17.66 ^{ab}	27.33 ^a	0.439 ^{ab}	0.0207 ^{bc}	3.33 ^a
200	541 ^a	11.00 ^{bc}	28.33 ^a	0.529 ^a	0.0223 ^{bc}	0.69 ^a
300	378 ^{ab}	12.66 ^{bc}	29.33 ^a	0.324 ^{bcd}	0.0290 ^{bc}	0.75 ^a
400	233 ^{bc}	18.33 ^{ab}	31.67 ^a	0.198 ^d	0.0323 ^{bc}	1.68 ^a
500	287 ^{bc}	25.00 ^a	31.67 ^a	0.239 ^{bcd}	0.0413 ^b	6.60 ^a
600	188 ^{bc}	11.00 ^{bc}	31.67 ^a	0.171 ^d	0.0483 ^b	1.37 ^a
700	170 ^c	7.66 ^c	32.67 ^a	0.217 ^{cd}	0.0553 ^b	7.58 ^a
800	271 ^{bc}	14.33 ^{bc}	35.00 ^a	0.258 ^{bcd}	0.0560 ^b	1.68 ^a
900	151 ^c	14.66 ^{bc}	35.00 ^a	0.250 ^{bcd}	0.1400 ^a	1.92 ^a
1000	154 ^c	11.66 ^{bc}	36.67 ^a	0.284 ^{bcd}	0.1413 ^a	1.93 ^a

Means followed by different letter(s) show significant difference at (P<0.05) significance level according to the Duncan's multiple ranges test.

Table 4. Inhibition index value (RI) of TiO₂ NPs different concentrations on seed Germination Percent (GP), Mean Germination Time (MGT), Germination Rate (GR), Germination Index (GI) and Weight Germination Index (WGI) of (*Vitex agnus-castus L.*)

TiO ₂ Concentration (ppm)	GP	MGT	GR	GI	WGI
10	-0.94	+0.06	-0.42	-0.47	-0.99
100	-0.01	+0.08	-0.06	+0.02	+0.03
200	+0.13	+0.03	+0.22	+0.22	+0.22
300	-0.24	+0.02	-0.38	-0.33	-0.32
400	-0.24	+0.02	-0.38	-0.31	-0.29
500	-0.24	+0.08	-0.12	-0.08	-0.09
600	-0.30	+0.02	-0.63	-0.62	-0.99
700	-0.30	+0.07	-0.38	-0.32	-0.31
800	-0.42	+0.03	-0.43	-0.41	-0.41
900	-0.50	+0.11	-0.47	-0.42	-0.42
1000	-0.50	+0.06	-0.55	-0.50	-0.99

(Table 3). The lowest optimal seedling index was observed at 10 µg.ml⁻¹ nitric oxide, which was lower than the control, and the highest optimum index was observed under 700 µg.ml⁻¹.

Titanium dioxide elongates plant organs such as root and shoot by increasing cell division (Farahani *et al.*, 2012). It seems that titanium dioxide nanoparticles act similarly to cytokinin and gibberellin and stimulate cell division and increase cell volume (Mandeh *et al.*, 2012). Gao *et al.* (2008) reported that titanium nanoparticles can enhance biomass growth in spinach leaves by increasing Rubisco enzyme activity. According to Khodakovskaya *et al.* (2009), titanium dioxide nanoparticles improved germination and root growth by increasing uptake by the thick shell of tomato seed and

increasing water uptake around the seed. Water absorption is important in seed germination because mature seeds are relatively dry and require significant amounts of water to initiate cell metabolism and seed growth. Determining the moisture content of seed and nano-materials inside the seeds supports this hypothesis; however, the mechanism of specific penetration through the seed shell and enhancing water uptake by nano-materials is not completely clear. Positive effects of titanium on plant growth such as cowpea by increasing number and length of pods and the increasing number of seeds per pod (Owolade *et al.*, 2008), increasing the fresh and dry weight of spinach seedlings by increasing nitrate uptake and leaf biomass (Mingyu *et al.*, 2007; Yang *et al.*, 2007), increasing seedling length and vigor of maize seed

RESEARCH ARTICLE

Table 5. Inhibition index value (RI) of TiO₂ NPs different concentrations on Seedling Vigor Index (SVI), Seedling Optimum Index (SOI), Radicle Length (RL), Plumule Length (PL), Wet Weight (WW) and Dry Weight (DW) on Seedling of (*Vitex agnus-castus* L.).

TiO ₂ Concentration (ppm)	SVI	SOI	RL	PL	WW	DW
10	0.03	-1.64	0.86	0.07	-0.17	0.42
100	0.44	0.54	0.79	0.10	0.06	0.96
200	0.65	-0.53	0.66	0.14	0.23	0.63
300	0.5	-0.50	0.71	0.17	-0.2	0.72
400	0.18	0.10	0.80	0.23	-0.52	0.75
500	0.34	0.77	0.85	0.23	-0.42	0.80
600	-0.005	-0.09	0.66	0.23	-0.57	0.83
700	-0.10	0.80	0.52	0.96	-0.47	0.85
800	0.30	0.10	0.74	0.30	-0.37	0.85
900	-0.20	0.21	0.75	0.30	-0.37	0.94
1000	-0.18	0.21	0.68	0.33	-0.3	0.94

RL (root length), PL (shoot length) and WW (fresh weight of seedling) and DW (dry weight of seedling).

(Buzea et al., 2007), increasing tomato seedling weight (Khodakovskaya et al., 2009), increasing fresh seedling weight of wheat (Mahmoodzadeh & Aghili, 2014), increasing fresh and dry weight of stem (Zhu et al., 2008), and an increasing number of apple leaves (Wojcik & Klamkowski, 2004) have been reported. Zhang et al. (2008) have reported that spraying titanium nanoparticles on cucumber leaves after several hours increased photosynthesis, which in turn increased root growth. Increased metabolism and growth of rice crop due to the use of Nano titanium dioxide have also been reported (Chutipaijit, 2015). In contrast, (Klancnik et al., 2011) reported that titanium dioxide nanoparticles did not affect the average root length and number of roots per tuber in onion. Thus, the uptake efficiency and the effect of nanoparticles on the growth and metabolic activities of different plants appear to be very different (Nair et al., 2010).

Response Index

Response index (RI) to different concentrations of titanium dioxide nanoparticles on germination and seedling growth indices are presented in (Tables 4 and 5).

The RI value is in the range of -1 and +1. Positive values indicate stimulation by treatment and negative values indicate inhibition compared to control. Titanium dioxide at all concentrations had a stimulating effect on mean germination time, root length, shoot length and dry weight (Tables 4 and 5). Also, titanium dioxide only had a stimulatory effect on germination percentage and germination rate in 200 mg.L⁻¹ treatment only. Treatments of 100 and 200 mg.L⁻¹ Nano titanium dioxide had a stimulating effect on the germination index and weighted germination index (Table 4). In general, titanium oxide nanoparticles had a stimulatory effect on most of the studied traits. However, its high concentrations also showed an inhibitory effect on some traits as shown in (Tables 4 and 5).

Conclusion

The Finally, it can be concluded that treatment with Nano titanium dioxide promoted germination and seedling growth, but this effect was not positive at all concentrations. Treatment of 200 µg.L⁻¹ titanium dioxide nanoparticles had the most stimulatory effect on germination and seedling growth factors of *Vitex* plantlets. The highest germination percentage, germination index, relative germination percentage, germination rate, germination index by weight, Vigor seedling index and seedling fresh weight were observed under 200 µg.ml⁻¹. As this concentration is relatively low, in addition to reducing the risk of adverse effects of the nanoparticles, it is also economically cost-effective and can be considered as a successful treatment to improve germination and early establishment of this valuable plant.

References

- Afzali A, Maghsoodlou SH. 2016. Modern application of nanotechnology in textile. - In: Nanostructured Polymer Blends and Composites in Textiles, (Apple Academic Press and CRC Press). pp.41-85.
- Aghdam MTB, Mohammadi H, Ghorbanpour M. 2016. Effects of Nano particulate anatase titanium dioxide on physiological and biochemical performance of (*Linum usitatissimum*) (Linaceae) under well-watered and drought stress conditions. *Braz J Bot*, 39(1): 139-146.
- Azadbakht M, Baheddini A, Shorideh SM, Naserzadeh A. 2005. Effect of (*Vitex agnus-castus* L.) leaf and fruit flavonoidal extracts on serum prolactin concentration. *J Med Plants*, 4(16):56-61.
- Azimi R, Jankju Borzelabad M, Feizi H, Azimi A. 2014. Interaction of SiO₂ nanoparticles with seed pre chilling on germination and

RESEARCH ARTICLE

- early seedling growth of tall wheatgrass (*Agropyron elongatum* L.). *Pol J Chem Technol*, 16(3):25-29.
- Buza C, Pacheco Blandino I, Robbie K. 2007. Nanomaterials and nanoparticles: sources and toxicity. *Bio interphases*, 2(4):17-71.
- Carmichael AR. 2008. Can Vitex (*Vitex agnus-castus* L.) be used for the treatment of mastalgia? What is the current evidence? *Evidence-Based Complementary and Alternative Medicine*, 5(3): 247-250.
- Chutipajit S. 2015. Establishment of condition and nanoparticle factors influencing plant regeneration from aromatic rice (*Oryza sativa*). *Int J Agric Biol*, 17(5): 1049-1054.
- Clement L, Charlot H, Marmier N. 2013. Toxicity of TiO₂ nanoparticle to cladocerans, algae rotifers and plants- Effects of size and crystalline structure. *Chemosphere*, 90(3):1083-1090.
- Du W, Sun Y, Ji R, Zhu J, Wu J., Guo H. 2011. TiO₂ and ZnO nanoparticles negatively affect wheat growth and soil enzyme activities in agricultural soil. *J Environ Monitor*, 13(4):822-828.
- Dugoua JJ, Seely D, Perri D, Koren G, Mills E. 2008. Safety and efficacy of chaste tree (*Vitex agnus-castus*) during pregnancy and lactation. *Journal of Population Therapeutics and Clinical Pharmacology*, 15(1). e74-e79.
- Elghniji K, Salem S, Mosbah MB, Elaloui E, Moussaoui Y. 2014. Detoxification of 4-chlorophenol in TiO₂ sunlight system: effect of raw and treated solution on seed germination and plants growth of various sensitive vegetables. *Toxicol Environ Chem*, 96(6): 869-879.
- Farahani HA, Sani B, Maroufi K. 2012. The germination variations in fleawort (*Plantago psyllium* L.) by nanoparticle. International conference on biotechnology, nanotechnology and its application. 17-18 March, Bangkok, Thailand. pp. 43-46.
- Fathi Z, Khavari Nejad RA, Mahmoodzadeh H, Nejad Satari T. 2017. Investigating of a wide range of concentrations of multi-walled carbon nanotubes on germination and growth of castor seeds (*Ricinus communis* L.). *J Plant Prod Res*, 57(3):228-236.
- Feizi H, Kamali M, Jafari L, Moghaddam PR. 2013. Phytotoxicity and stimulatory impacts of nanosized and bulk titanium dioxide on fennel (*Foeniculum vulgare* Mill). *Chemosphere*, 91(4):506-511.
- Frazier TP, Burklew CE, Zhang B. 2014. Titanium dioxide nanoparticles affect the growth and microRNA expression of tobacco (*Nicotiana tabacum*). *Functional & Integrative Genomics*, 14(1):75-83.
- Gao F, Liu C, Qu C, Zheng L, Yang F, Su M, Hong F. 2008. Was improvement of spinach growth by nano-TiO₂ treatment related to the changes of Rubisco activase? *Biomaterials*, 21(2):211-217.
- Hatami M, Ghorbanpour M, Salehjarjomand H. 2014. Nano-anatase TiO₂ modulates the germination behavior and seedling vigority of some commercially important medicinal and aromatic plants. *J Biol Environ Sci*, 8(22):53-59.
- Jaberzadeh A, Moaveni P, Moghadam HRT, Zahedi H. 2013. Influence of bulk and nanoparticles titanium foliar application on some agronomic traits, seed gluten and starch contents of wheat subjected to water deficit stress. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 41(1):201-207.
- Kandil AA, Sharief AE, Ahmed KR. 2015. Performance of some soybean (*Glycine max* L.) Merrill. cultivars under salinity stress to germination characters. *Int J Agrar Agricul Res*, 6(3):48-56.
- Kenanakis G, Katsarakis N. 2014. Chemically grown TiO₂ on glass with superior photocatalytic properties. *J Environ Chem Eng*, 2(3):1748-1755.
- Khodakovskaya M, Dervishi E, Mahmood M, Xu Y, Li Z, Watanabe F, Biris AS. 2009. Carbon nanotubes are able to penetrate plant seed coat and dramatically affect seed germination and plant growth. *Am Chem Soc*, 3(10):3221-3227.
- Klancnik K, Drobne D, Valant J, Dolenc Koce J. 2011. Use of a modified Allium test with nanoTiO₂. *Ecotox Environ Saf*.74(1):85-92.
- Kulkarni MG, Street RA, Van Staden J. 2007. Germination and seedling growth requirements for propagation of *Dioscorea dregeana* (Kunth) Dur. and Schinz—a tuberous medicinal plant. *South Afr J Bot*, 73(1):131-137.
- Laware SL, Raskar SV. 2014. Effect of titanium dioxide nanoparticles on hydrolytic and antioxidant enzymes during seed germination in onion. *Int J Curr Microbiol Appl Sci*, 3(7):749-760.
- Lu CM, Zhang CY, Wen JQ, Wu GR, Tao, MX. 2002. Research of the effect of nanometer materials on germination and growth enhancement of (*Glycine max*) and its mechanism. *Soya Bean Science*, 21(3):168-171.
- Mahajan P, Dhoke SK, Khanna AS. 2011. Effect of nano-ZnO particle suspension on growth of mung (*Vigna radiata*) and gram (*Cicer arietinum*) seedlings using plant agar method. *J Nanotechnol*. Volume 2011, Article ID 696535, 7 pages.
- Mahmoodzadeh H, Aghili R. 2014. Effect on germination and early growth characteristics in wheat plants (*Triticum aestivum* L.) seeds exposed to TiO₂ nanoparticles. *Journal of Chemical Health Risks*, 4(1):29-36.
- Mandeh M, Omidi M, Rahaie M. 2012. In Vitro Influences of TiO₂ Nanoparticles on Barley (*Hordeum vulgare* L.) Tissue Culture. *Biological Trace Element Research*, 150(1-3): 376-380.
- Marschner H. 2011. Marschner's Mineral Nutrition of Higher Plants. 3rd edition. Edited by P. Marschner. Amsterdam, Netherlands: Elsevier/Academic Press, pp. 684. ISBN 978-0-12-384905-2.
- Mingyu S, Hong F, Liu C, Wu X, Liu X, Chen L, Gao F, Yang F, Li Z. 2007. Effects of Nano-anatase TiO₂ on absorption, distribution of light and photo reduction activities of chloroplast membrane of Spinach. *Biol Trace Elem Res*, 120(1-3):273-283.
- Nair PMG, Chung IM. 2014. A mechanistic study on the toxic effect of copper oxide nanoparticles in soybean (*Glycine max* L.) root development and lignification of root cells. *Biol Trace Elem Res*, 162(1-3):342-352.
- Nair R, Varghese SH, Nair BG, Maekawa T, Yoshida Y, Kumar DS. 2010. Nano particulate material delivery to plants. *Plant Sci*, 179(3):154-163.
- Owolade OF, Ogunleti DO, Adenekan MO. 2008. Effects of Titanium dioxide on the diseases, development and yield of edible cowpea. *Electron J Environ Agricul Food Chem*, 7(5):2942-2947.
- Pais I. 2008. The biological importance of titanium. *J Plant Nutr*, 6(1):3-131.
- Panwar P, Bhardwaj SD. 2005. Handbook of practical forestry. Agrobios (India). p 191.
- Qi M, Liu Y, Li T. 2013. Nano-TiO₂ improve the photosynthesis of tomato leaves under mild heat stress. *Biol Trace Elem Res*, 156(1-3):323-328.
- Rani A, Sharma A. 2013. The genus Vitex: A review. *Pharmacognosy reviews*, 7(14): p.188.
- Rico CM, Peralta-Videa JR, Gardea-Torresdey JL. 2015. Differential effects of cerium oxide nanoparticles on rice, wheat, and barley roots: A Fourier Transform Infrared (FT-IR) micro spectroscopy study. *Appl Spectrosc*, 69(2):287-295.
- Siddiqui MH, Al-Whaibi MH, Firoz M, Al-Khaishany MY. 2015. Role of nanoparticles in plants. - In *Nanotechnology and Plant Sciences* (pp. 19-35). Springer, Cham.
- Syu YY, Hung JH, Chen JC, Chuang HW. 2014. Impacts of size and shape of silver nanoparticles on Arabidopsis plant growth and gene expression. *Plant Physiol Biochem*, 83, pp.57-64.
- Tiwari PK, Singh AK, Singh VP, Prasad SM, Ramawat N, Tripathi DK, Chauhan DK, Rai AK. 2019. Liquid assisted pulsed laser ablation synthesized copper oxide nanoparticles (CuO-NPs) and their differential impact on rice seedlings. *Ecotoxic Environ Saf*, 176, pp.321-329.
- Tripathi DK, Singh S, Singh S, Pandey R, Singh VP, Sharma NC, Prasad SM, Dubey NK, Chauhan DK. 2017. An overview on

RESEARCH ARTICLE

- manufactured nanoparticles in plants: uptake, translocation, accumulation and phytotoxicity. *Plant Physiol Biochem*, 110, pp.2-12.
- Tripathi DK, Singh S, Singh VP, Prasad SM, Chauhan DK, Dubey NK. 2016. Silicon nanoparticles more efficiently alleviate arsenate toxicity than silicon in maize cultivar and hybrid differing in arsenate tolerance. *Front Environ Sci*, 4, p.46.
- Vashisth A, Nagarajan S. 2010. Effect on germination and early growth characteristics in sunflower (*Helianthus annuus*) seeds exposed to static magnetic field. *J Plant Physiol*, 167(2):149-156.
- Wojcik P, Klamkowski K. 2004. "Szampion" apple tree response to foliar titanium application. *J Plant Nutr*, 27(11):2033-2046.
- Wu G, Du, G. 2007. Germination is related to seed mass in grasses (Poaceae) of the eastern Qinghai-Tibetan Plateau, China. *Nordic J Bot*, 25(5-6):361-365.
- Yang L, Watts DJ. 2005. Particle surface characteristics may play an important role in phytotoxicity of alumina nanoparticles. *Toxicol Lett*, 158(2):122-132.
- Yang Y, Liu Q, Han C, Qiao YZ, Yao XQ, Yin HJ. 2007. Influence of water stress and low irradiance on morphological and physiological characteristics of (*Picea asperata*) seedlings. *Photosynthetica*, 45(4):613-619.
- Zhang P, Cui HX, Zhang ZJ, Zhong RG. 2008. Effects of nano-TiO₂ photo semiconductor on photosynthesis of cucumber plants. *Chinese Agricultural Science Bulletin*, 24 pp.230-233.
- Zheng L, Hong F, Lu S, Liu C. 2005. Effect of nano-TiO₂ on strength of naturally aged seeds and growth of spinach. *Biol Trace Elem Res*, 104(1):83-91.
- Zhu H, Han J, Xiao JQ, Jin Y. 2008. Uptake, translocation and accumulation of manufactured iron oxide nanoparticles by pumpkin plants. *J Environ Monitor*, 10(6):713-717.